		Birth	1 Month	2 Months	4 Months	6 Months
Re	Hepatitis B	□ Нер В	☐ Hep B¹			□ Нер В
Recommended Immunizations	Rotavirus			□RV	□RV	□RV
	Diphtheria, Tetanus, Pertussis			□DTaP	□ DТаР	□DTaP
	Haemophilus influenzae type b			□Hib	□Hib	□Hib
	Pneumococcal			□PCV	□PCV	□PCV
	Inactivated Poliovirus			□IPV	□IPV	□IPV
tions	Influenza					☐ Influenza, first dose²☐ second dose
Milestones	Milestones should be achieved by the end of the age indicated. Talk to your healthcare provider about ageappropriate milestones if your child was born prematurely.	Recognizes caregiver's voice	Responsive to calming actions when upset Starts to smile Raises head when on tummy	 □ Developing a social smile □ Raises chest when lying on tummy □ Follows objects with eyes □ Makes cooing noises 	☐ Grasps and shakes hand toys ☐ Laughs out loud ☐ Babbles more expressively ☐ Holds head steady	Responds to own name Reaches with one hand Recognizes familiar faces Rolls over
Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	WEIGHT / PERCENTILE LENGTH / PERCENTILE	WEIGHT / PERCENTILE LENGTH / PERCENTILE	WEIGHT / PERCENTILE LENGTH / PERCENTILE	WEIGHT / PERCENTILE LENGTH / PERCENTILE	WEIGHT / PERCENTILE LENGTH / PERCENTILE
		HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE	HEAD CIRCUMFERENCE
	Shaded boxes indicate the vaccine can be given during shown age range.		VISIT DATE	VISIT DATE	VISIT DATE	VISIT DATE

² Children 6 months or older should receive flu vaccination every flu season. If this is the first time for flu vaccine, a child 6 months through 8 years of age should receive two doses, separated by at least 4 weeks. If this child only receives one dose in the first season, he or she should receive two doses the next season, if still younger than 9 years. Ask your child's healthcare provider if a second dose is needed.



1-800-CDC-INFO (1-800-232-4636) or visit

http://www.cdc.gov/vaccines/recs/schedules/child-schedule (Immunization)

http://www.cdc.gov/actearly/growthchart (Milestones)

¹ The second dose of Hep B may be given either at the 1 month or 2 month visit.

		12 MONTHS	15 MONTHS	18 MONTHS	19-23 MONTHS	2-3 YEARS	4-6 YEARS
	Hepatitis B	☐ Hep B (Final dose admi	nistered between 6 and 18 r	months)			
Recommended Immunizations	Diphtheria, Tetanus, Pertussis		□DTaP				□DTaP
	Haemophilus influenzae type b	□Hib					
	Pneumococcal	□ PCV ²					
	Inactivated Poliovirus	□IPV					□IPV
	Influenza	☐ Influenza, first dose ☐ second dose (if needed)	,3			Age Age 2 3 Influenza, first dose³ second dose (if needed)	Age Age Age 4 5 6 Influenza, first dose ³ second dose (if needed)
	Measles, Mumps, Rubella	□MMR					□MMR
	Varicella	□ Varicella					□Varicella
	Hepatitis A	☐☐ Hep A (2 doses be	etween 6 and 23 months				
Milestones	Milestones should be achieved by the end of the age indicated. Talk to your healthcare provider about ageappropriate milestones if your child was born prematurely.	☐ Cries when mother or father leaves ☐ Finger-feeds himself ☐ Explores objects in many different ways (shaking, banging, throwing, dropping) ☐ Says "dada" and "mama"	☐ Drinks from a cup☐ Scribbles☐ Indicates what he wants by pointing	☐ Says at least 6 words with meaning ☐ Points to one body part ☐ Walks up steps	☐ Completes simple chores when asked ☐ Is spontaneous with affection	☐ Points to object or picture when named ☐ Uses 2- to 4-word phrases ☐ Follows simple instructions ☐ Gradually becomes more interested in other children	☐ Balances on one foot ☐ Shows school readiness skills ☐ Prints some letters and numbers ☐ Has mature pencil grasp ☐ Has good articulation/ language skills
Growth	At each well child visit, enter date, length, weight, and percentile information to keep track of your child's progress.	WEIGHT / PERCENTILE LENGTH / PERCENTILE HEAD CIRCUMFERENCE	WEIGHT / PERCENTILE LENGTH / PERCENTILE HEAD CIRCUMFERENCE	WEIGHT / PERCENTILE LENGTH / PERCENTILE HEAD CIRCUMFERENCE	WEIGHT / PERCENTILE LENGTH / PERCENTILE HEAD CIRCUMFERENCE	WEIGHT HEIGHT BMI	WEIGHT HEIGHT BMI
	Shaded boxes indicate the vaccine can be given during shown age range.	VISIT DATE	VISIT DATE	VISIT DATE	VISIT DATE	VISIT DATE	VISIT DATE

Immunizations and Developmental Milestones for Your Child from Birth through 6 years old

³ Children 6 months or older should receive flu vaccination every flu season. If this is the first time for flu vaccine, a child 6 months through 8 years of age should receive two doses, separated by at least 4 weeks. If this child only receives one dose in the first season, he or she should receive two doses the next season, if still younger than 9 years. Ask your child's healthcare provider if a second dose is needed.



Child's Name

1-800-CDC-INFO (1-800-232-4636) or visit

http://www.cdc.gov/vaccines/recs/schedules/child-schedule (Immunization)

Birth Date

http://www.cdc.gov/actearly/growthchart (Milestones)

The Recommended Immunization Schedule for persons aged birth through 6 years old is approved by the Centers for Disease Control and Prevention, the American Academy of Pediatrics, and the American Academy of Family Physicians

² HepA vaccination is recommended for high-risk children older than 2 years, along with a dose of meningococcal vaccine (MCV4) and pneumococcal vaccine (PPSV). HepA vaccination may be administered to any child over 2 for whom immunity is desired. See vaccine-specific recommendations at http://www.cdc.gov/vaccines/pubs/ACIP-list.htm.